



Gersteflocken

Pasze jednoskładnikowe

Feed material

Tak jak w przypadku kukurydzy, zalety jęczmienia mogą zostać w pełni wykorzystane tylko kiedy poddamy go obróbce hydrotermicznej. Płatkowany jęczmień może znaleźć zastosowanie jako wartościowa podstawa dla Twojej własnej indywidualnej mieszanki paszowej albo jako wzbogacenie zwykłej paszy treściwej. "Czasem mniej znaczy więcej" - zastąp 1 kg owsa 0.9 kg jęczmienia, aby osiągnąć porównywalny poziom energii.

Przejdź do sklepu: <https://eggersmann.com.pl/pasze-jednoskladnikowe/54-gersteflocken.html>

Dawkowanie:

Feeding recommendation:

for horses:

- Barley was traditionally used as an energy source in horse feeding in the Orient and is now often used in the feed rations of sport and breeding horses.
- As an alternative to oats, the higher energy content must be taken into account!
- 1kg of oats is replaced with 0.9kg of barley flakes.

for chicken & other poultry:

- Barley flakes are used in chicken feed to increase energy levels, but due to the high content of β -glucans, the rate at which they are mixed into the complete feed should be limited in order to ensure good usability of the feed
- For pullets and chicken chicks we recommend a mixing rate of approx. 15-20%, for laying hens, laying quail and parent animals up to 40%, parent animals of waterfowl up to 60%.

for small animals:

- As with all types of grain, care should be taken with barley so as not to burden the digestive tract of rabbits and rodents.
- In special situations, however, it can make sense to energetically upgrade the feeding.
- Our barley flakes can, for example, be offered over the winter months or for pregnant animals as an energy-rich feed supplement.
- To keep the animals busy, we recommend distributing 0.5-1 teaspoon under the fresh bedding or in the hay.

białko og. strawne (vRp): 83,4 g/kg
preceacal digestible protein (pcvRp): 67,9 g/kg
energia strawna (MJ DE): 12,5 MJ DE/kg
Metabolizable energy (MJ ME): 11,6 MJ ME/kg





Analiza: 10,60 % białko surowe, 2,00 % tłuszcz surowy, 5,00 % włókno surowe, 2,50 % popiół surowy, 0,07 % wapń, 0,35 % fosfor, 52,8 % skrobia, 2,30 % cukier

